

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5

BIVALENT ANTI-PSMGFR STIMULATES CELL GROWTH IN MUC1⁺ BREAST TUMOR CELLS - 1504

1504 CELLS (6000 PER WELL) STIMULATED EVERY 24 OR 48 HOURS WITH ANTI-PSMGFR AND HARVESTED AFTER 5 DAYS (---) OR 6 DAYS (----)

**Fig. 6**




Fig. 7

ANTI-PSMGFR STIMULATES GROWTH OF BREAST TUMOR CELL LINE 1500

1500 CELLS STIMULATED EVERY 24h OR 48h WITH ANTI-PSMGFR AND HARVESTED AT 2 DAYS (---), 3 DAYS (==), OR 4 DAYS (----)




Fig. 8

BIVALENT ANTI-PSMGFR STIMULATES CELL PROLIFERATION IN T47D CELLS
THE EFFECT OF α -PSMGFR BIVALENT ANTIBODY ON T47D CELL PROLIFERATION




Fig. 9




Fig. 10

MONOVALENT ANTI-PSMGFR INHIBITS GROWTH OF BREAST TUMOR CELL LINE 1504




Fig. 11

MONOVALENT ANTI-PSMGFR INHIBITS GROWTH OF BREAST TUMOR CELL LINE 1500

EFFECT OF MONOVALENT ANTI-PSMGFR (24hr INTERVAL) ON 1500
CELL PROLIFERATION - HARVESTED AFTER 7 DAYS




Fig. 12




Fig. 13




Fig. 14




Fig. 15

BIVALENT ANTI-PSMGFR INDUCES MAP KINASE CELL
PROLIFERATION PATHWAY IN 1504 BREAST TUMOR CELLS

TIME-COURSE EFFECT OF ANTI-PSMGFR (5 μ l) ON ERK2 PATHWAY IN 1504 CELL LINE




Fig. 16

BIVALENT ANTI-PSMGFR ALSO INDUCES ERK2
PHOSPHORYLATION IN 1500 BREAST TUMOR CELLS

TIME-COURSE EFFECT OF ANTI-PSMGFR (5uI) ON ERK2 PATHWAY IN 1500 CELL LINE




Fig. 17

**MN COMPOUNDS COMPETE WITH ANTIBODY TO BIND TO PSMGFR &
BLOCK ACTIVATION OF MAP KINASE PROLIFERATION PATHWAY**




Fig. 18

MONOVALENT INHIBITS ERK2 PHOSPHORYLATION
ANTI-PSMGFR (5ul) AND MONOVALENT (10ul) ANTIBODY EFFECT
ON ERK2 PATHWAY IN 1500 CELL LINE




Fig. 19

BREAST TUMOR CELLS PRESENT FULL-LENGTH AS WELL AS CLEAVED MUC1




Fig. 20




Fig. 21




Fig. 22




Fig. 23

MONOVALENT ANTI-PSMGFR INHIBITS CELL GROWTH IN NAT-PSMGFRTG TRANSFECTANTS

BIVALENT ANTI-PSMGFR (■) OR MONOVALENT (□)
(1uI) & BIVALENT NAT-PSMGFRTG TRANSFECTANT




Fig. 24




Fig. 25

IN NAT-PSMGFRTC TRANSFECTANTS BIVALENT (5ul)
ACTIVATES ERK2 - MONOVALENT (10ul) INHIBITS




Fig. 26




Fig. 27




Fig. 28

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.